

TRAFFIC SAFETY FACTS Crash • Stats

DOT HS 811 851

A Brief Statistical Summary

November 2013

Lives Saved in 2012 by Restraint Use and **Minimum Drinking Age Laws**

In 2012, the use of seat belts in passenger vehicles saved an estimated 12,174 lives. Seat belts have saved nearly 63,000 lives during the 5-year-period from 2008 to 2012. These annual estimates of lives saved, shown in Table 1, have been produced by NHTSA's National Center for Statistics and Analysis since 1975. The estimates are calculated using the effectiveness of each device or law that is mentioned. More information on the methodology of lives-saved estimates is available in two NHTSA publications, Lives Saved FAQs (NHTSA, 2009, DOT HS 811 105), which answers 30 common questions about lives saved and includes references to many other reports on lives saved; and Lives Saved Calculations for Seat Belts and Frontal Air Bags (Glassbrenner & Starnes, 2009, DOT HS 811 206, NHTSA), which describes in detail the methodology of estimating lives saved by seat belts and frontal air bags.

In addition to the 12,174 lives saved in 2012 by seat belts (occupants 5 and older), 2,213 lives were saved by frontal air bags (occupants 13 and older), 1,699 lives were saved by motorcycle helmets, 525 lives were saved by 21-yearold-minimum-drinking-age laws, and 284 lives (4 and younger) were saved by child restraints (child safety seats and lap/shoulder belts). An additional 3,031 lives would have been saved in 2012 if all unrestrained passenger vehicle occupants 5 and older involved in fatal crashes had worn their seat belts; and if all motorcyclists had been helmeted, then an additional 781 lives would have been saved.

Table 1 shows that the number of lives saved by seat belts among occupants 5 and older increased from 11,983 in 2011 to 12,174 in 2012. The number of lives saved by child restraints, frontal air bags, and motorcycle helmets also increased from 2011 to 2012. The number of lives saved by the minimum-drinking-age law decreased from 535 in 2011 to 525 in 2012.

	Lives Saved, Age 4 & Younger	Lives Saved, Age 5 & Older	Lives Saved, Age 13 & Older	Lives Saved, All Ages	Lives Saved	Additional Lives That Would Have Been Saved at 100 Percent Use	
Year	Child Restraints	Seat Belts	Frontal Air Bags	Motorcycle Helmets	Minimum Drinking Age Law	Seat Belts	Motorcycle Helmets
2008	286	13,312	2,557	1,836	716	4,171	827
2009	307	12,763	2,387	1.486	626	3,700	733
2010	303	12,582	2,315	1,556	552	3,353	708
2011	262	11,983	2,210	1,622	535	3,394	706
2012	284	12,174	2,213	1,699	525	3,031	781

Table 1: Lives Saved by Restraint Use, and 21-Year-Old-Minimum-Drinking-Age Laws, and Additional Lives That Would Have Been Saved at 100-Percent Seat Belt and Motorcycle Helmet Use, 2008–2012

Source: 2008–2011 FARS Final Files and FARS 2012 Annual Report File

Table 2 shows lives-saved totals in 2012 for all 50 States, the District of Columbia, and Puerto Rico. The far left column is the lives saved by child restraints, and the next three columns refer respectively to lives saved by seat belts, lives saved by frontal air bags, and lives saved by motor-

cycle helmets. The numbers in the two far right columns represent the additional lives savable at 100-percent seat belt use and 100-percent helmet use, respectively. Due to rounding, the sum of the State totals may not equal the national total.

	Lives Saved by	Lives Saved by	Lives Saved by	Lives Saved by	Additional Lives Savable At 100-Percent Restraint Use	
State	Child Restraints, Age 4 and Younger	Seat Belts, Age 5 and Older	Frontal Air Bags, Age 13 and Older	Motorcycle Helmets, All Ages	Seat Belts	Motorcycle Helmets
Alabama	11	304	69	52	75	3
Alaska	0	21	3	3	5	2
Arizona	4	231	45	41	90	27
Arkansas	3	189	42	15	108	17
California	26	1,194	140	239	96	10
Colorado	2	149	30	15	58	20
Connecticut	0	90	16	7	22	10
Delaware	1	35	7	8	7	2
District of Columbia	0	0	0	2	0	0
Florida	12	752	129	135	190	98
Georgia	15	486	85	74	85	3
Hawaii	0	27	4	7	4	11
Idaho	0	73	12	7	31	4
Illinois	12	346	63	17	47	45
				17	47	45
Indiana	13	316	53 26			
lowa	5	161		7	26	18
Kansas	5	143	31	7	61	13
Kentucky	6	272	57	23	97	26
Louisiana	1	246	48	45	101	1
Maine	1	55	15	6	20	5
Maryland	3	209	33	41	35	3
Massachusetts	1	95	23	29	50	1
Michigan	11	417	66	42	50	25
Minnesota	3	173	28	8	21	16
Mississippi	9	203	50	20	82	2
Missouri	5	195	67	56	111	4
Montana	0	48	16	6	34	8
Nebraska	2	62	18	12	31	0
Nevada	2	90	15	21	17	2
New Hampshire	0	25	8	6	19	7
New Jersey	1	181	32	40	43	3
New Mexico	2	132	20	14	26	15
New York	7	460	61	91	79	6
North Carolina	10	507	79	105	119	7
North Dakota	1	55	13	2	26	4
Ohio	9	379	89	22	148	47
Oklahoma	1	263	53	12	95	24
Oregon	0	146	16	28	10	2
Pennsylvania	10	328	102	62	136	39
Rhode Island	1	32	5	1	13	2
South Carolina	4	278	61	25	61	39
South Dakota	2	40	9	23	31	8
Tennessee	7	373	77	79	133	2
Texas	50	1,479	214	107	185	101
Utah Vermont	4 2	<u> </u>	15 6	12 5	27	4
Vermont					7	1
Virginia	2	276	57	46	120	2
Washington	2	176	23	46	12	2
West Virginia	2	91	28	10	39	5
Wisconsin	4	230	45	16	83	34
Wyoming	1	55	9	1	24	4
National*	284	12,174	2,213	1,699	3,031	781
Puerto Rico	0	61	15	10	15	13

Table 2: Lives Saved in 2012 by Child Restraints, Seat Belts, Frontal Air Bags, and Motorcycle Helmets

*Estimates for States may not add up to national total due to independent rounding.